PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau

INTERNATIONAL AP

ATION PUBLISHED UNDER THE PATE

OOPERATION TREATY (PCT)

(51) International Patent Classification 5:

G01N 33/58, 21/55

(11) International Publication Number:

WO 91/02981

A1

(43) International Publication Date:

7 March 1991 (07.03.91)

(21) International Application Number:

PCT/GB90/01319

(22) International Filing Date:

24 August 1990 (24.08.90)

(30) Priority data:

8919411.2

25 August 1989 (25.08.89)

GB

(71) Applicant (for all designated States except US): AMER-SHAM INTERNATIONAL PLC [GB/GB]; Amersham Place, Little Chalfont, Buckinghamshire HP7 9NA (GB).

(72) Inventors; and (75) Inventors/Applicants (for US only): GARLAND, Peter, Bryan [GB/GB]; Hope Cottage, Sunnyway, Old Bosham, West Sussex PO18 8HQ (GB). CHARLES, Stephen, Alexander [GB/GB]; 5 Eliot Close, Heydon Hill, Aylesbury, Buckinghamshire HP1 9JB (GB).

(74) Agent: PENNANT, Pyers; Stevens, Hewlett & Perkins, 5 Quality Court, Chancery Lane, London WC2A 1HZ

(81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent)*, DK (European patent), ES (European patent), FR (European patent), GB (European patent), IT (European pa pean patent), JP, LU (European patent), NL (European patent), SE (European patent), US.

Published

With international search report.

(54) Title: ASSAY METHOD

(57) Abstract

A method of assaying for an analyte by the use of surface plasmon resonance spectrometry (SPRS). The analyte is a member of a non-immune ligand-receptor pair. A metal surface carries the analyte or an analogue immobilised with the other member of the ligand-receptor pair reversibly bound thereto. A fluid containing the analyte is brought into contact with the metal surface, and displacement of the other member of the ligand-receptor pair monitored by SPRS.

^{*} See back of page

DESIGNATIONS OF "DE"

Until further notice, any designation of "DE" in any international application whose international filing date is prior to October 3, 1990, shall have effect in the territory of the Federal Republic of Germany with the exception of the territory of the former German Democratic Republic.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

ΑT	Austria	ES	Spain	MC	Monaco
AU	Australia	FI	Finland	MG	Madagascar
BB	Barbados	FR	France	ML	Mali
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Fasso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GR	Greece	NL	Netherlands
BJ	Benin	HU	Hungary	NO	Norway
BR	Brazil	IT	Italy	PL	Poland
CA	Canada	9L	Japan	RO	Romania
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CC	Congo		of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CM	Cameroon .	LI	Liechtenstein	รข	Soviet Union
DΕ	Germany	LK	Sri Lanka	TD	Chad .
DK	Denmark	ŁU	Luxembourg	TG	Togo
				US	United States of America

WO 91/02981 PCT/GB90/01319

5

15

35

metal.

ASSAY METHOD

This invention concerns a method of assaying for a macromolecular analyte by use of the technique of surface plasmon resonance spectrometry (SPRS).

The phenomenon of SPR is well known and will not be described in detail (see EPA 305109 for example). Briefly, the intensity of monochromatic plain-polarised light (conveniently obtained from a laser) reflected from the interface between an optically transparent material, e.g. glass, and a thin metal layer depends on the refractive index of material on the downstream side of the metal. Accordingly, by measuring changes in intensity of the reflected light an indication can be obtained of changes in refractive index of material at a particular point on the down-stream surface of the

The intensity of reflected light also varies with the angle of incidence, and reflectivity drops 20 sharply to a minimum at a particular angle which is characteristic of the equipment.

WO 89/08260 lescribes a method of analysing for an analyte in a sample by bringing the sample into contact with a metal surface, on which an antibody has 25 previously been reversibly bound to immobilised analyte or analogue, and monitoring displacement of antibody as indicative of the presence or the concentration of the analyte in the sample. specification is mainly concerned with hapten analytes 30 and describes only hapten-antibody and antigenantibody binding pairs.

EPA 276142 describes competition assays involving an analyte and two other reagents, in which SPRS is used to monitor the formation of a complex on a solid surface. The assay involves the use of at

5

10

20

Growth factor or

analogue

least one other liquid reagent in addition to the sample. The signal resulting from complex formation on the surface may be obscured by noise resulting from non-specific binding of other macromolecules to the surface.

This invention provides a method of assaying for an analyte, preferably a macromolecular analyte, which is a member of a ligand-receptor pair other than a hapten-antibody or an antigen-antibody pair, by the use of a metal surface adapted for surface plasmon resonance spectrometry which metal surface carries the analyte or an analogue thereof immobilised thereon with the other member of the ligand-receptor pair reversibly bound thereto, which method comprises bringing a fluid containing the analyte into contact with the metal surface and observing by surface plasmon resonance spectrometry displacement of the other member of the ligand-receptor pair from the The analyte is a member of a ligandreceptor pair. Many examples of such pairs are known and include the following:

	Ligand	Receptor
	DNA	DNA
	DNA	RNA
25	RNA	RNA
	Protein	DNA
	Protein	RNA
	DNA	Binding drug
	RNA	Binding drug
30	Lectin	Oligosaccharide (free or in a
		glycoprotein or glycolipid)
	Neurotransmitter or	
	analogue	Protein receptor
	Hormone	Hormone receptor

Protein receptor

35

- 3 -

Differentiation factor

or analogue Protein receptor

Enzyme Cofactor, substrate or inhibitor

Biotin Avidin

5 Enzyme Prosthetic group

Oligopeptide Protein
Immunoglobulin Protein A

Drug Binding protein

In the above, the term protein is used to include peptides.

The method involves the use of a metal surface adapted for surface plasmon resonance spectrometry. The metal may comprise silver or gold, conveniently in the form of a layer e.g. deposited by evaporation on a 15 carrier such as a glass slide. The metal surface carries the analyte or an analogue thereof immobilised An analogue of the analyte is a substance which competes with the analyte for binding to the other member of the ligand-receptor pair. 20 analogue will be arranged to be as near as possible or even completely identical to the analyte. The use in assays of analyte analogues is well known. of the analyte or analogue to the metal surface without loss of binding power is effected by methods 25 which are well known. Particularly when the analyte or analogue is of low molecular weight, the use of a spacer molecule may be required. To prevent nonspecific binding at a later stage in the assay, any surplus area of the metal surface may be coated e.g. 30 with an inert protein.

Reversibly bound to the immobilised analyte or analogue is the other member of the ligand-receptor pair. In another aspect, this invention provides an assay device comprising a metal surface adapted for surface plasmon resonance spectrometry, which surface

10

carries immobilised thereon a member of a ligandreceptor pair, other than a hapten-antibody or an
antigen-antibody pair, with the other member of the
ligand-receptor pair reversibly bound thereto. That
other member of the ligand-receptor pair may have been
modified to increase the signal generated to increase
the SPR signal generated by its removal from the metal
surface. Such modification may involve addition of
a molecule or group of low refractive index, or more
preferably high refractive index such as polystyrene,
titanium dioxide or gold colloid.

Using this pre-formed device, the method of the invention is very simple. A fluid containing the analyte is brought into contact with the pre-coated metal surface. Analyte in the fluid sample competes with immobilised analyte for binding to the other member of the ligand-receptor pair. Displacement of the other member of the ligand-receptor pair into solution, as a complex with analyte in the sample, is monitored by surface plasmon resonance spectrometry. The method has two particular advantages:

- (a) The only fluid reagent involved is the sample containing the analyte. When this is brought into contact with the coated metal surface, the presence or the concentration of the analyte in the sample can be assayed within minutes or even seconds.
- the SPRS signal, generated by removal of a reagent from the metal surface, is not significantly contaminated by noise due to non-specific binding of material to the surface.

25

30

- 5 -

Example 1

Displacement Assay using Nucleic Acid Probes

a complementary oligonucleotide B (17mer).

Oligonucleotide B was then covalently linked to the silver surface of a slide. The slide was then blocked with hybridisation buffer. A further oligonucleotide C (50mer) which had the same sequence as part of oligonucleotide B and therefore complementary to A, was then added and displacement from the silver slide of the hybridised oligonucleotide A was measured. Results using 32P end-labelled oligonucleotide A showed that as the amount of oligonucleotide C was increased more oligonucleotide A was released. Such changes will also be measurable by SPR.

Example 2

20 DNA Strand Displacement by SPR

Method

Preparation of oligonucleotides

A sixteen mer probe and a complementary ninety-seven mer target were prepared using phosphoramidite chemistry on the Applied Biosystems Model 380D DNA synthesizer. The DNA probe was modified in order to permit efficient and stable binding to the lver surface. This was prepared by attaching a terminal primary amine at the 5´-end of the molecule. The amino-oligonucleotide was mixed with an equal volume of 0.1M sodium hydrogen carbonate solution ph 8.5 and to this mixture a solution of SPDP (0.25 mg for each 1.0 OD of oligo used) in DMF was added. The reaction was left to equilibriate for 90 minutes at

room temperature and then eluted through a Sephadex G25 PD10 column. Fractions containing the required product were pooled together and purified by preparative HPLC. After purification the appropriate fractions were dried down by vacuum centrifugation to remove HPLC solvents and then reconstituted in a known volume of water.

Preparation of DNA Hybrid

and a complementary 97 mer was prepared by mixing 16 mer and 97 mer in a 1.5:1 molar ratio ensuring that the 16 mer, the immobilisable probe, was in excess. The preannealing was carried out at 65°C in 2x SSPE (300mM NaC1, 20mM NaH₂PO₄.H₂O, 2mM ethylenediaminetetra-acetic acid) and the mixture was allowed to cool down to room temperature for 2hrs.

SPR Experiment

- Using 1.8 x 10^{-10} moles of hybrid/ml, 1ml was pumped across a silver slide at 1µl/s after priming the slide with 2x SSPE. Following a wash step with 1ml of 2x SSPE the slide was blocked with hybridisation buffer followed by a 2x SSPE wash.
- 3×10^{-10} moles of a 50 mer, which is complementary in sequence to the 97 mer, in 1ml 2x SSPE was flowed across the slide at a speed of 1µl/s, and the change in reflectivity was monitored with time.
- Control reactions involving no DNA and non-30 complementary DNA were carried out.

Data is presented below with the No DNA control value deducted from the experimental.

- 7 -

Table 1

	Target DNA	<u>-6</u>	change	<u>1 n</u>	reflectivity
5	No DNA control				
	NO DNA CONTROL				0
	50 mer			-	- 1
	Non-complementary	DNA		+	· 1

Results demonstrate displacement of the
97 mer sequence away from the silver surface by the
complementary 50 base sequence. This does not occur in
the absence of this oligonucleotide nor in the presence
of the non-complementary sequence which does not
hybridise to the 97 mer.

20

25

30

35

Į.

- 8 -

CLAIMS

5


1. A method of assaying for an analyte which is a member of a ligand-receptor pair other than a hapten-antibody or an antigen-antibody pair, by the use of a metal surface adapted for surface plasmon resonance spectrometry which metal surface carries the analyte or an analogue thereof immobilised thereon with the other member of the ligand-receptor pair reversibly bound thereto, which method comprises bringing a fluid containing the analyte into contact with the metal surface and observing by surface plasmon resonance spectrometry displacement of the other member of the ligand-receptor pair from the surface.

- 2. A method as claimed in claim 1, wherein the analyte is a macromolecule.
- 3. An assay device comprising a metal surface adapted for surface plasmon resonance spectrometry, which surface carries immobilised thereon a member of a ligand-receptor pair, other than a hapten-antibody or an antigen-antibody pair, with the other member of the ligand-receptor pair reversibly bound thereto.

30

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶							
According to International Patent Classification (IPC) or to both National Classification and IPC							
IPC5: G 01 N 33/58, 21/55							
II. FIELDS SEARCHED							
Minimum Documentation Searched ⁷							
Classification System Classification Symbols							
IPC5	IPC5 G 01 N						
	٠	Documentation Searched oth to the Extent that such Docume	er than Minimum Documentation nts are included in Fields Searched ^a				
		DNSIDERED TO BE RELEVANT					
Category •		on of Document, ¹¹ with Indication, where a		Relevant to Claim No.13			
X .	se	, 276142 (ARES-SERONO RESMITED PARTNERSHIP) 27 Ju e page 1, line 1; page 3 ble 1; cited in the appl	ly 1988, . line 20	1-3			
X	se pa	8703093 (RADIOMETER A/S) e page 6, line 34; page 7 ge 9, line 1 - line 18; aims 1,7-9) 21 May 1987, 7, line 20;	1-3			
Ρ,Χ	se pa	8909408 (ARES-SERONO RES MITED PARTNERSHIP) 5 Octo e page 5, line 18; page 6 ge 8, line 8 - line 20 ble 2, claims	ber 1989.	1-3			
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date of priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention							
"L" doci	filing date "L" document which may throw doubts on gricrity claim(s) or which is cited to establish the publication of solution of the considered to the co						
Of document referring to an oral disclosure, use, exhibition or other means P document published prior to the international discount of the art.							
V. CERTIFICATION							
Date of the Actual Completion of the International Search Date of Mailing of this International Search Report 1. 12. 90							
Internationa	I Searching	Authority	Signature of Authorized Officer				
	EUROPEAN PATENT OFFICE miss T. MORTENSEN THE STATE OF THE						

Form PCT/ISA/210 (second sheet) (January 1985)



Form PCT/ISA/210 (extra sheet) (January 1985)

AN TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/GB 90/01319

SA

39697

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent ffice EDP file on 27/09/90

The European Patent office is in no way liable for theseparticulars which are merely given for the purpose of information.

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
EP-A0-	276142	27/07/88	NONE		
WO-A-	8703093	21/05/87	EP-A-	0245396	19/11/87
WO-A-	8909408	05/10/89	AU-D- EP-A-	3361889 0359807	16/10/89 28/03/90
US÷A- 	4626513	02/12/86	WO-A-	88/04429	16/06/88
EP-A-	0245206	11/11/87	AU-D- JP-T- WO-A-	7583887 1500221 87/06956	01/12/87 26/01/89 19/11/87
WO-A-	8908260	08/09/89	AU-D- EP-A-	3077489 0378594	31/08/89 25/07/90